# Computer Vision

# Homogeneous Coordinates

Julien SEINTURIER
*Associate Professor*

julien.seinturier@univ-tln.fr
http://web.seinturier.fr/teaching/computer-vision

# Representing 3D space

3D space can be represented with Euclidean Space

- Vector space on $\mathbb{R}^3$
- Orthonormal basis of 3 vectors denoted X, Y and Z
- dot product $\cdot$ and cross product $\times$



# Representing 3D space

3D space can be represented with Euclidean Space

- Vector space on $\mathbb{R}^3$
- Orthonormal basis of 3 vectors denoted X, Y and Z
- dot product $\cdot$ and cross product $\times$

Mathematical properties

- Euclidean norm $\|u\| = \sqrt{u \cdot u}$
- Angle: $\cos\theta = \frac{u \cdot v}{\|u\|\|v\|}$



# Representing Transforms

| Transform | Computation | Linearity | Distances | Angles |
|---|---|---|---|---|
| Translation | $(\alpha + x, \beta + y, \gamma + z)$ | Affine | ✓ | ✓ |
| Rotation | $\begin{pmatrix} x \\ y\cos(\omega) - z\sin(\omega) \\ y\sin(\omega) + z\cos(\omega) \end{pmatrix}$ $\begin{pmatrix} x\cos(\varphi) + z\sin(\varphi) \\ y \\ z\cos(\varphi) - x\sin(\varphi) \end{pmatrix}$ $\begin{pmatrix} x\cos(\kappa) - y\sin(\kappa) \\ x\sin(\kappa) + y\cos(\kappa) \\ z \end{pmatrix}$ | Linear | ✓ | ✓ |
| Scale (uniform) | $(sx, sy, sz)$ | Linear | ✗ | ✓ |
| Scale | $(s_x x, s_y y, s_z z)$ | Linear | ✗ | ✗ |

# Representing Transforms

Complex computation

| Transform | Computation | Linearity | Distances | Angles |
|---|---|---|---|---|
| Translation | $(\alpha + x, \beta + y, \gamma + z)$ | Affine | ✓ | ✓ |
| Rotation | $\begin{pmatrix} x \\ y\cos(\omega) - z\sin(\omega) \\ y\sin(\omega) + z\cos(\omega) \end{pmatrix}$ $\begin{pmatrix} x\cos(\varphi) + z\sin(\varphi) \\ y \\ z\cos(\varphi) - x\sin(\varphi) \end{pmatrix}$ $\begin{pmatrix} x\cos(\kappa) - y\sin(\kappa) \\ x\sin(\kappa) + y\cos(\kappa) \\ z \end{pmatrix}$ | Linear | ✓ | ✓ |
| Scale (uniform) | $(sx, sy, sz)$ | Linear | ✗ | ✓ |
| Scale | $(s_x x, s_y y, s_z z)$ | Linear | ✗ | ✗ |

# Representing Transforms

Various properties

| Transform | Computation | Linearity | Distances | Angles |
|---|---|---|---|---|
| Translation | $(\alpha + x, \beta + y, \gamma + z)$ | Affine | ✓ | ✓ |
| Rotation | $\begin{pmatrix} x \\ y\cos(\omega) - z\sin(\omega) \\ y\sin(\omega) + z\cos(\omega) \end{pmatrix}$ $\begin{pmatrix} x\cos(\varphi) + z\sin(\varphi) \\ y \\ z\cos(\varphi) - x\sin(\varphi) \end{pmatrix}$ $\begin{pmatrix} x\cos(\kappa) - y\sin(\kappa) \\ x\sin(\kappa) + y\cos(\kappa) \\ z \end{pmatrix}$ | Linear | ✓ | ✓ |
| Scale (uniform) | $(sx, sy, sz)$ | Linear | ✗ | ✓ |
| Scale | $(s_x x, s_y y, s_z z)$ | Linear | ✗ | ✗ |

# Representing 3D space

- Limits
  - No uniform representation of transformations
  - Computation can be complex
- Needs
  - A consistent mathematical formalism
  - Good computational properties

## Definition[Equivalence relation]

*Let $E$ be a set. An **equivalence relation** on $E$, denoted $\sim$, is a binary operation that satisfies:*

## Definition[Equivalence relation]

*Let $E$ be a set. An **equivalence relation** on $E$, denoted $\sim$, is a binary operation that satisfies:*

- $\forall a \in E, a \sim a$ **Reflexivity**

## Definition[Equivalence relation]

*Let $E$ be a set. An **equivalence relation** on $E$, denoted $\sim$, is a binary operation that satisfies:*

- $\forall a \in E, a \sim a$ **Reflexivity**
- $\forall a, b \in E, a \sim b \leftrightarrow b \sim a$ **Symmetry**

## Definition[Equivalence relation]

*Let $E$ be a set. An **equivalence relation** on $E$, denoted $\sim$, is a binary operation that satisfies:*

- $\forall a \in E, a \sim a$ **Reflexivity**
- $\forall a, b \in E, a \sim b \longleftrightarrow b \sim a$ **Symmetry**
- $\forall a, b, c \in E, a \sim b \; and \; b \sim c \longrightarrow a \sim c$ **Transitivity**

## Definition[Equivalence relation]

*Let $E$ be a set. An **equivalence relation** on $E$, denoted $\sim$, is a binary operation that satisfies:*

- $\forall a \in E, a \sim a$ **Reflexivity**
- $\forall a, b \in E, a \sim b \longleftrightarrow b \sim a$ **Symmetry**
- $\forall a, b, c \in E, a \sim b \; and \; b \sim c \longrightarrow a \sim c$ **Transitivity**

## Definition[Equivalence class]

*Let $E$ be a set with an equivalence relation $\sim$ and let $x \in E$. The **equivalence class** of $x$ for $\sim$ is the subset, denoted $\bar{x}^{\sim}$ such as:*

$$\bar{x}^{\sim} = \{y \in E, x \sim y\}$$

## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation

  A line $x$ is parallel to itself





## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation

  A line $x$ is parallel to itself

  If a line $x$ is parallel to a line $y$, then $y$ is parallel to $x$






# Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation

  A line $x$ is parallel to itself

  If a line $x$ is parallel to a line $y$, then $y$ is parallel to $x$

  If a line $x$ is parallel to a line $y$ and if $y$ is parallel to $z$ then $x$ is parallel to $z$





## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation
- The subset $G = \{a, b, c, d, e\}$ is an equivalent class



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation
- The subset $G = \{a, b, c, d, e\}$ is an equivalent class
- The subset $H = \{f, g, h, i\}$ is an equivalent class



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation
- The subset $G = \{a, b, c, d, e\}$ is an equivalent class
- The subset $H = \{f, g, h, i\}$ is an equivalent class
- $G \cap H = \emptyset$



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation
- The subset $G = \{a, b, c, d, e\}$ is an equivalent class
- The subset $H = \{f, g, h, i\}$ is an equivalent class
- $G \cap H = \emptyset$
- $G \cup H = E$



## Example[Equivalence]

Let $E = \{a, b, c, d, e, f, g, h, i\}$ be a set of lines within the plane and let $\sim$ a binary operation defined such as $\forall x, y \in E, x \sim y \rightarrow x\ parallel\ to\ y$

- $\sim$ is an equivalence relation
- The subset $G = \{a, b, c, d, e\}$ is an equivalent class
- The subset $H = \{f, g, h, i\}$ is an equivalent class
- $G \cap H = \emptyset$
- $G \cup H = E$

Equivalent classes form a partition



## Definition[Projective space]

*Let $E$ be a vector space on a field $\mathbb{K}$. The projective space $P(E)$ is the set of equivalence classes of $E \setminus \{0_E\}$ under an equivalence relation, denoted $\sim$, such as:*

$$\forall \mathcal{X}, \mathcal{Y} \in E \setminus \{0_E\},\ \mathcal{X} \sim \mathcal{Y} \text{ if } \exists \lambda \in \mathbb{K},\ \lambda \neq 0, \mathcal{X} = \lambda \mathcal{Y}$$

## Definition[Projective space]

*Let $E$ be a vector space on a field $\mathbb{K}$. The projective space $P(E)$ is the set of equivalence classes of $E \setminus \{0_E\}$ under an equivalence relation, denoted $\sim$, such as:*

$$\forall \mathcal{X}, \mathcal{Y} \in E \setminus \{0_E\},\ \mathcal{X} \sim \mathcal{Y} \text{ if } \exists \lambda \in \mathbb{K},\ \lambda \neq 0, \mathcal{X} = \lambda \mathcal{Y}$$

## Property[Projective space dimension]

Let $E^n$ be a vector space of finite dimension $n$. The projective space $P(E^n)$ has a dimension of $n - 1$.

## Definition[Homogeneous vector]

*A **homogeneous vector**, denoted $\mathcal{X}$, is defined such that:*

$$\forall \lambda \in \mathbb{R}, \lambda \neq 0, \mathcal{X} = \lambda \mathcal{X}$$

## Definition[Homogeneous vector]

*A **homogeneous vector**, denoted $\mathcal{X}$, is defined such that:*

$$\forall \lambda \in \mathbb{R}, \lambda \neq 0, \mathcal{X} = \lambda \mathcal{X}$$

## Interpretation

- A homogeneous vector describes an equivalence class of collinear vectors

## Definition[Homogeneous vector]

*A **homogeneous vector**, denoted $\mathcal{X}$, is defined such that:*

$$\forall \lambda \in \mathbb{R}, \lambda \neq 0, \mathcal{X} = \lambda \mathcal{X}$$

## Interpretation

- A homogeneous vector describes an equivalence class of collinear vectors
- A homogenous vector has an infinity of representations:

$$\chi = (1,4,6) = \left(\frac{1}{2}, 2,3\right) = (12,48,72) = \cdots = (\lambda \times 1, \lambda \times 4, \lambda \times 6)$$

$\lambda = 1$ $\lambda = 1/2$ $\lambda = 12$

## Homogeneous Representation

- **Definition**: A homogeneous vector $\mathcal{X}$ of dimension $n$ is a tuple made of $n$ **homogenous coordinates**:

$$\forall \lambda \in \mathbb{R}^* \quad \mathcal{X} = \lambda(x_1, \ldots, x_n) = (\lambda x_1, \ldots, \lambda x_n)$$

## Homogeneous Representation

- **Definition**: A homogeneous vector $\mathcal{X}$ of dimension $n$ is a tuple made of $n$ **homogenous coordinates**:

$$\forall \lambda \in \mathbb{R}^* \quad \mathcal{X} = \lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

- A homogeneous vector $\mathcal{X} = \lambda(x_1, \dots, x_n)$ can be represented by a column matrix, denoted X, such as:

$$\mathrm{X} = \lambda \begin{bmatrix} x_1 \\ \cdots \\ x_n \end{bmatrix}$$

## Homogeneous Representation

- **Definition**: A homogeneous vector $\mathcal{X}$ of dimension $n$ is a tuple made of $n$ **homogenous coordinates**:

$$\forall \lambda \in \mathbb{R}^* \quad \mathcal{X} = \lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

- A homogeneous vector $\mathcal{X} = \lambda(x_1, \dots, x_n)$ can be represented by a column matrix, denoted X, such as:

$$\mathrm{X} = \lambda \begin{bmatrix} x_1 \\ \cdots \\ x_n \end{bmatrix} = \begin{bmatrix} \lambda x_1 \\ \cdots \\ \lambda x_n \end{bmatrix}$$

Column notation

## Homogeneous Representation

- **Definition**: A homogeneous vector $\mathcal{X}$ of dimension $n$ is a tuple made of $n$ **homogenous coordinates**:

$$\forall \lambda \in \mathbb{R}^* \quad \mathcal{X} = \lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

- A homogeneous vector $\mathcal{X} = \lambda(x_1, \dots, x_n)$ can be represented by a column matrix, denoted X, such as:

$$\mathrm{X} = \lambda \begin{bmatrix} x_1 \\ \cdots \\ x_n \end{bmatrix} = \begin{bmatrix} \lambda x_1 \\ \cdots \\ \lambda x_n \end{bmatrix} = [\lambda x_1 \quad \cdots \quad \lambda x_n]^\top = \lambda [x_1 \quad \cdots \quad x_n]^\top$$

Column notation — Row notation

## Euclidean to homogeneous Representation

- **Definition:** Let $a = (a_1, \dots, a_n)$ be a vector within a Euclidean space of dimension $n$. A **homogeneous representation** of $a$ is a vector of dimension $n + 1$, denoted $\mathcal{A}$, such as:

$$\mathcal{A} = (\lambda a_1, \dots, \lambda a_n, \lambda) \equiv A = \begin{bmatrix} \lambda a_1 \\ \dots \\ \lambda a_n \\ \lambda \end{bmatrix} = [\lambda a_1 \quad \dots \quad \lambda a_n \quad \lambda]^\top$$

## Euclidean to homogeneous Representation

- **Definition:** Let $a = (a_1, \dots, a_n)$ be a vector within a Euclidean space of dimension $n$. A **homogeneous representation** of $a$ is a vector of dimension $n + 1$, denoted $\mathcal{A}$, such as:

$$\mathcal{A} = (\lambda a_1, \dots, \lambda a_n, \lambda) \equiv A = \begin{bmatrix} \lambda a_1 \\ \dots \\ \lambda a_n \\ \lambda \end{bmatrix} = [\lambda a_1 \quad \dots \quad \lambda a_n \quad \lambda]^\top$$

- For the sake of simplicity, $\lambda$ is usually set to $1$

# Euclidean to homogeneous Representation

- **Example:** From Euclidean to homogeneous

$$(1, 4, 3)$$

Euclidean vector

# Euclidean to homogeneous Representation

- **Example:** From Euclidean to homogeneous

$$(1,4,3) \rightarrow [1,4,3,1]^{\mathsf{T}} = 2[1,4,3,1]^{\mathsf{T}} = \cdots = \lambda[1,4,3,1]^{\mathsf{T}}$$

Euclidean vector

Homogeneous representations

# Euclidean to homogeneous Representation

- **Example:** From Euclidean to homogeneous

$$(1,4,3) \rightarrow [1,4,3,1]^{\mathsf{T}} = 2[1,4,3,1]^{\mathsf{T}} = \cdots = \lambda[1,4,3,1]^{\mathsf{T}}$$

Euclidean vector

Homogeneous representations

- **Warning:** Homogeneous representation of the Euclidean space origin **is not** the origin of homogeneous space:

$$(0,0,0) \rightarrow [0,0,0,1]^{\mathsf{T}} = 2[0,0,0,1]^{\mathsf{T}} = \cdots = \lambda[0,0,0,1]^{\mathsf{T}}$$

## Homogeneous to Euclidean representation

- Let $\mathcal{A} = [a_1 \quad \dots \quad a_n \quad w]^\top$ be a vector within a homogeneous space of dimension $n+1$, with $w \neq 0$.

  The Euclidean representation of $\mathcal{A}$ is a vector of dimension $n$, denoted $A$ such as:

$$A = \left(\frac{a_1}{w}, \dots, \frac{a_n}{w}\right)$$

# Homogeneous to Euclidean representation

- **Example:** From homogeneous to Euclidean

$$[2 \quad 8 \quad 6 \quad 2]^T$$

homogeneous vector

# Homogeneous to Euclidean representation

- **Example:** From homogeneous to Euclidean

$$[2 \quad 8 \quad 6 \quad 2]^{\top} \rightarrow \left(\frac{2}{2}, \frac{8}{2}, \frac{6}{2}\right) = (1,4,3)$$

homogeneous vector

Euclidean representation

# Homogeneous to Euclidean representation

- **Example:** From homogeneous to Euclidean

$$[2 \quad 8 \quad 6 \quad 2]^{\top} \rightarrow \left(\frac{2}{2}, \frac{8}{2}, \frac{6}{2}\right) = (1,4,3)$$

homogeneous vector — Euclidean representation

- **Warning:** Homogeneous space origin **cannot be represented** within Euclidean space:

$$[0,0,0,0]^{\top} \rightarrow \left(\frac{0}{0}, \frac{0}{0}, \frac{0}{0}\right)$$

# Geometric interpretation

- Let $A = (x, y)$ a vector within a 2D Euclidean space



# Geometric interpretation

- Let $A = (x, y)$ a vector within a 2D Euclidean space
- $\mathcal{A} = [x \quad y \quad 1]^\top$ represents $A$ within the homogeneous space



# Geometric interpretation

- Let $A = (x, y)$ a vector within a 2D Euclidean space
- $\mathcal{A} = [x \quad y \quad 1]^\top$ represents $A$ within the homogeneous space
- $\lambda\mathcal{A} = [\lambda x \quad \lambda y \quad \lambda]^\top$ is another representation of $A$



# Geometric interpretation

- Let $A = (x, y)$ a vector within a 2D Euclidean space
- $\mathcal{A} = [x \quad y \quad 1]^\top$ represents $A$ within the homogeneous space
- $\lambda\mathcal{A} = [\lambda x \quad \lambda y \quad \lambda]^\top$ is another representation of $A$




All homogeneous representations of $A$ are lying on the same line

## Homogeneous transformation

- Let be a homogeneous space of dimension $n$ and let $\mathcal{V}$ be a homogeneous vector. A **homogeneous transform** is an application defined such as:

$$\mathcal{H}(\mathcal{V}) = \mathrm{HV}$$

With:

- $\mathrm{H}$ is a $n \times n$ square matrix that represents $\mathcal{H}$
- $V$ is a $n$ dimensioned vector that represents $\mathcal{V}$
- $\mathrm{HV}$ is the matrix product of $\mathrm{H}$ and $\mathrm{V}$

# Homogeneous transformation

- Within 4 dimensioned homogeneous space
  - $\mathcal{V} = [x \quad y \quad z \quad w]^{\top}$
  - H is such as:

$$\mathrm{H} = \begin{bmatrix} h_{11} & h_{12} & h_{13} & h_{14} \\ h_{21} & h_{22} & h_{23} & h_{24} \\ h_{31} & h_{32} & h_{33} & h_{34} \\ h_{41} & h_{42} & h_{43} & h_{44} \end{bmatrix}$$

# Homogeneous transformation

- Identity transform

$$\mathrm{H} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Computation

$$\mathrm{H}\mathcal{V} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity
  - Let $\mathcal{H}$ be a homogeneous transform represented by matrix $\mathrm{H}$:

$$H = \begin{bmatrix} h_{11} & h_{12} & h_{13} & h_{14} \\ h_{21} & h_{22} & h_{23} & h_{24} \\ h_{31} & h_{32} & h_{33} & h_{34} \\ h_{41} & h_{42} & h_{43} & h_{44} \end{bmatrix}$$

  - Let $\mathcal{A}$ and $\mathcal{B}$ two homogeneous vectors represented by column matrices $\mathrm{A}$ and $\mathrm{B}$ respectively and let $\lambda$ and $\mu$ be two scalars:

$$\mathrm{A} = \begin{bmatrix} x_a \\ y_a \\ z_a \\ w_a \end{bmatrix} \quad \mathrm{B} = \begin{bmatrix} x_b \\ y_b \\ z_b \\ w_b \end{bmatrix} \quad \lambda\mathrm{A} = \begin{bmatrix} \lambda x_a \\ \lambda y_a \\ \lambda z_a \\ \lambda w_a \end{bmatrix} \quad \mu\mathrm{B} = \begin{bmatrix} \mu x_b \\ \mu y_b \\ \mu z_b \\ \mu w_b \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A} + \mu\mathcal{B}) = \mathrm{H}(\lambda\mathrm{A} + \mu\mathrm{B})$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A} + \mu\mathcal{B}) = \mathrm{H}(\lambda\mathrm{A} + \mu\mathrm{B})$$

$$\mathcal{H}(\lambda\mathcal{A} + \mu\mathcal{B}) = \begin{bmatrix} h_{11} & h_{12} & h_{13} & h_{14} \\ h_{21} & h_{22} & h_{23} & h_{24} \\ h_{31} & h_{32} & h_{33} & h_{34} \\ h_{41} & h_{42} & h_{43} & h_{44} \end{bmatrix} \begin{bmatrix} \lambda x_a + \mu x_b \\ \lambda y_a + \mu y_b \\ \lambda z_a + \mu z_b \\ \lambda w_a + \mu w_b \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \mathrm{H}(\lambda\mathcal{A}+\mu\mathcal{B})$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11} & h_{12} & h_{13} & h_{14} \\ h_{21} & h_{22} & h_{23} & h_{24} \\ h_{31} & h_{32} & h_{33} & h_{34} \\ h_{41} & h_{42} & h_{43} & h_{44} \end{bmatrix} \begin{bmatrix} \lambda x_a + \mu x_b \\ \lambda y_a + \mu y_b \\ \lambda z_a + \mu z_b \\ \lambda w_a + \mu w_b \end{bmatrix}$$

Matrix product

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}(\lambda x_a + \mu x_b) + h_{12}(\lambda y_a + \mu y_b) + h_{13}(\lambda z_a + \mu z_b) + h_{14}(\lambda w_a + \mu w_b) \\ h_{21}(\lambda x_a + \mu x_b) + h_{22}(\lambda y_a + \mu y_b) + h_{23}(\lambda z_a + \mu z_b) + h_{24}(\lambda w_a + \mu w_b) \\ h_{31}(\lambda x_a + \mu x_b) + h_{32}(\lambda y_a + \mu y_b) + h_{33}(\lambda z_a + \mu z_b) + h_{44}(\lambda w_a + \mu w_b) \\ h_{41}(\lambda x_a + \mu x_b) + h_{42}(\lambda y_a + \mu y_b) + h_{43}(\lambda z_a + \mu z_b) + h_{44}(\lambda w_a + \mu w_b) \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11} & h_{12} & h_{13} & h_{14} \\ h_{21} & h_{22} & h_{23} & h_{24} \\ h_{31} & h_{32} & h_{33} & h_{34} \\ h_{41} & h_{42} & h_{43} & h_{44} \end{bmatrix} \begin{bmatrix} \lambda x_a + \mu x_b \\ \lambda y_a + \mu y_b \\ \lambda z_a + \mu z_b \\ \lambda w_a + \mu w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}(\lambda x_a + \mu x_b) + h_{12}(\lambda y_a + \mu y_b) + h_{13}(\lambda z_a + \mu z_b) + h_{14}(\lambda w_a + \mu w_b) \\ h_{21}(\lambda x_a + \mu x_b) + h_{22}(\lambda y_a + \mu y_b) + h_{23}(\lambda z_a + \mu z_b) + h_{24}(\lambda w_a + \mu w_b) \\ h_{31}(\lambda x_a + \mu x_b) + h_{32}(\lambda y_a + \mu y_b) + h_{33}(\lambda z_a + \mu z_b) + h_{44}(\lambda w_a + \mu w_b) \\ h_{41}(\lambda x_a + \mu x_b) + h_{42}(\lambda y_a + \mu y_b) + h_{43}(\lambda z_a + \mu z_b) + h_{44}(\lambda w_a + \mu w_b) \end{bmatrix}$$

Distribution

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{11}\mu x_b + h_{12}\lambda y_a + h_{12}\mu y_b + h_{13}\lambda z_a + h_{13}\mu z_b + h_{14}\lambda w_a + h_{14}\mu w_b \\ h_{21}\lambda x_a + h_{21}\mu x_b + h_{22}\lambda y_a + h_{22}\mu y_b + h_{23}\lambda z_a + h_{23}\mu z_b + h_{24}\lambda w_a + h_{24}\mu w_b \\ h_{31}\lambda x_a + h_{31}\mu x_b + h_{32}\lambda y_a + h_{32}\mu y_b + h_{33}\lambda z_a + h_{33}\mu z_b + h_{44}\lambda w_a + h_{34}\mu w_b \\ h_{41}\lambda x_a + h_{31}\mu x_b + h_{42}\lambda y_a + h_{42}\mu y_b + h_{43}\lambda z_a + h_{43}\mu z_b + h_{44}\lambda w_a + h_{44}\mu w_b \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}(\lambda x_a+\mu x_b)+h_{12}(\lambda y_a+\mu y_b)+h_{13}(\lambda z_a+\mu z_b)+h_{14}(\lambda w_a+\mu w_b) \\ h_{21}(\lambda x_a+\mu x_b)+h_{22}(\lambda y_a+\mu y_b)+h_{23}(\lambda z_a+\mu z_b)+h_{24}(\lambda w_a+\mu w_b) \\ h_{31}(\lambda x_a+\mu x_b)+h_{32}(\lambda y_a+\mu y_b)+h_{33}(\lambda z_a+\mu z_b)+h_{44}(\lambda w_a+\mu w_b) \\ h_{41}(\lambda x_a+\mu x_b)+h_{42}(\lambda y_a+\mu y_b)+h_{43}(\lambda z_a+\mu z_b)+h_{44}(\lambda w_a+\mu w_b) \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a+h_{11}\mu x_b+h_{12}\lambda y_a+h_{12}\mu y_b+h_{13}\lambda z_a+h_{13}\mu z_b+h_{14}\lambda w_a+h_{14}\mu w_b \\ h_{21}\lambda x_a+h_{21}\mu x_b+h_{22}\lambda y_a+h_{22}\mu y_b+h_{23}\lambda z_a+h_{23}\mu z_b+h_{24}\lambda w_a+h_{24}\mu w_b \\ h_{31}\lambda x_a+h_{31}\mu x_b+h_{32}\lambda y_a+h_{32}\mu y_b+h_{33}\lambda z_a+h_{33}\mu z_b+h_{44}\lambda w_a+h_{34}\mu w_b \\ h_{41}\lambda x_a+h_{31}\mu x_b+h_{42}\lambda y_a+h_{42}\mu y_b+h_{43}\lambda z_a+h_{43}\mu z_b+h_{44}\lambda w_a+h_{44}\mu w_b \end{bmatrix}$$

Grouping

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a+h_{12}\lambda y_a+h_{13}\lambda z_a+h_{14}\lambda w_a+h_{11}\mu x_b+h_{12}\mu y_b+h_{13}\mu z_b+h_{14}\mu w_b \\ h_{21}\lambda x_a+h_{22}\lambda y_a+h_{23}\lambda z_a+h_{24}\lambda w_a+h_{21}\mu x_b+h_{22}\mu y_b+h_{23}\mu z_b+h_{24}\mu w_b \\ h_{31}\lambda x_a+h_{32}\lambda y_a+h_{33}\lambda z_a+h_{34}\lambda w_a+h_{31}\mu x_b+h_{32}\mu y_b+h_{33}\mu z_b+h_{34}\mu w_b \\ h_{41}\lambda x_a+h_{42}\lambda y_a+h_{43}\lambda z_a+h_{44}\lambda w_a+h_{41}\mu x_b+h_{42}\mu y_b+h_{43}\mu z_b+h_{44}\mu w_b \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{11}\mu x_b + h_{12}\lambda y_a + h_{12}\mu y_b + h_{13}\lambda z_a + h_{13}\mu z_b + h_{14}\lambda w_a + h_{14}\mu w_b \\ h_{21}\lambda x_a + h_{21}\mu x_b + h_{22}\lambda y_a + h_{22}\mu y_b + h_{23}\lambda z_a + h_{23}\mu z_b + h_{24}\lambda w_a + h_{24}\mu w_b \\ h_{31}\lambda x_a + h_{31}\mu x_b + h_{32}\lambda y_a + h_{32}\mu y_b + h_{33}\lambda z_a + h_{33}\mu z_b + h_{44}\lambda w_a + h_{34}\mu w_b \\ h_{41}\lambda x_a + h_{31}\mu x_b + h_{42}\lambda y_a + h_{42}\mu y_b + h_{43}\lambda z_a + h_{43}\mu z_b + h_{44}\lambda w_a + h_{44}\mu w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{12}\lambda y_a + h_{13}\lambda z_a + h_{14}\lambda w_a + h_{11}\mu x_b + h_{12}\mu y_b + h_{13}\mu z_b + h_{14}\mu w_b \\ h_{21}\lambda x_a + h_{22}\lambda y_a + h_{23}\lambda z_a + h_{24}\lambda w_a + h_{21}\mu x_b + h_{22}\mu y_b + h_{23}\mu z_b + h_{24}\mu w_b \\ h_{31}\lambda x_a + h_{32}\lambda y_a + h_{33}\lambda z_a + h_{34}\lambda w_a + h_{31}\mu x_b + h_{32}\mu y_b + h_{33}\mu z_b + h_{34}\mu w_b \\ h_{41}\lambda x_a + h_{42}\lambda y_a + h_{43}\lambda z_a + h_{44}\lambda w_a + h_{41}\mu x_b + h_{42}\mu y_b + h_{43}\mu z_b + h_{44}\mu w_b \end{bmatrix}$$

Decomposition

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{12}\lambda y_a + h_{13}\lambda z_a + h_{14}\lambda w_a \\ h_{21}\lambda x_a + h_{22}\lambda y_a + h_{23}\lambda z_a + h_{24}\lambda w_a \\ h_{31}\lambda x_a + h_{32}\lambda y_a + h_{33}\lambda z_a + h_{34}\lambda w_a \\ h_{41}\lambda x_a + h_{42}\lambda y_a + h_{43}\lambda z_a + h_{44}\lambda w_a \end{bmatrix} + \begin{bmatrix} h_{11}\mu x_b + h_{12}\mu y_b + h_{13}\mu z_b + h_{14}\mu w_b \\ h_{21}\mu x_b + h_{22}\mu y_b + h_{23}\mu z_b + h_{24}\mu w_b \\ h_{31}\mu x_b + h_{32}\mu y_b + h_{33}\mu z_b + h_{34}\mu w_b \\ h_{41}\mu x_b + h_{42}\mu y_b + h_{43}\mu z_b + h_{44}\mu w_b \end{bmatrix}$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{12}\lambda y_a + h_{13}\lambda z_a + h_{14}\lambda w_a + h_{11}\mu x_b + h_{12}\mu y_b + h_{13}\mu z_b + h_{14}\mu w_b \\ h_{21}\lambda x_a + h_{22}\lambda y_a + h_{23}\lambda z_a + h_{24}\lambda w_a + h_{21}\mu x_b + h_{22}\mu y_b + h_{23}\mu z_b + h_{24}\mu w_b \\ h_{31}\lambda x_a + h_{32}\lambda y_a + h_{33}\lambda z_a + h_{34}\lambda w_a + h_{31}\mu x_b + h_{32}\mu y_b + h_{33}\mu z_b + h_{34}\mu w_b \\ h_{41}\lambda x_a + h_{42}\lambda y_a + h_{43}\lambda z_a + h_{44}\lambda w_a + h_{41}\mu x_b + h_{42}\mu y_b + h_{43}\mu z_b + h_{44}\mu w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{12}\lambda y_a + h_{13}\lambda z_a + h_{14}\lambda w_a \\ h_{21}\lambda x_a + h_{22}\lambda y_a + h_{23}\lambda z_a + h_{24}\lambda w_a \\ h_{31}\lambda x_a + h_{32}\lambda y_a + h_{33}\lambda z_a + h_{34}\lambda w_a \\ h_{41}\lambda x_a + h_{42}\lambda y_a + h_{43}\lambda z_a + h_{44}\lambda w_a \end{bmatrix} + \begin{bmatrix} h_{11}\mu x_b + h_{12}\mu y_b + h_{13}\mu z_b + h_{14}\mu w_b \\ h_{21}\mu x_b + h_{22}\mu y_b + h_{23}\mu z_b + h_{24}\mu w_b \\ h_{31}\mu x_b + h_{32}\mu y_b + h_{33}\mu z_b + h_{34}\mu w_b \\ h_{41}\mu x_b + h_{42}\mu y_b + h_{43}\mu z_b + h_{44}\mu w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda \begin{bmatrix} h_{11}x_a + h_{12}y_a + h_{13}z_a + h_{14}w_a \\ h_{21}x_a + h_{22}y_a + h_{23}z_a + h_{24}w_a \\ h_{31}x_a + h_{32}y_a + h_{33}z_a + h_{34}w_a \\ h_{41}x_a + h_{42}y_a + h_{43}z_a + h_{44}w_a \end{bmatrix} + \mu \begin{bmatrix} h_{11}x_b + h_{12}y_b + h_{13}z_b + h_{14}w_b \\ h_{21}x_b + h_{22}y_b + h_{23}z_b + h_{24}w_b \\ h_{31}x_b + h_{32}y_b + h_{33}z_b + h_{34}w_b \\ h_{41}x_b + h_{42}y_b + h_{43}z_b + h_{44}w_b \end{bmatrix}$$

Factorization

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \begin{bmatrix} h_{11}\lambda x_a + h_{12}\lambda y_a + h_{13}\lambda z_a + h_{14}\lambda w_a \\ h_{21}\lambda x_a + h_{22}\lambda y_a + h_{23}\lambda z_a + h_{24}\lambda w_a \\ h_{31}\lambda x_a + h_{32}\lambda y_a + h_{33}\lambda z_a + h_{34}\lambda w_a \\ h_{41}\lambda x_a + h_{42}\lambda y_a + h_{43}\lambda z_a + h_{44}\lambda w_a \end{bmatrix} + \begin{bmatrix} h_{11}\mu x_b + h_{12}\mu y_b + h_{13}\mu z_b + h_{14}\mu w_b \\ h_{21}\mu x_b + h_{22}\mu y_b + h_{23}\mu z_b + h_{24}\mu w_b \\ h_{31}\mu x_b + h_{32}\mu y_b + h_{33}\mu z_b + h_{34}\mu w_b \\ h_{41}\mu x_b + h_{42}\mu y_b + h_{43}\mu z_b + h_{44}\mu w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda \begin{bmatrix} h_{11}x_a + h_{12}y_a + h_{13}z_a + h_{14}w_a \\ h_{21}x_a + h_{22}y_a + h_{23}z_a + h_{24}w_a \\ h_{31}x_a + h_{32}y_a + h_{33}z_a + h_{34}w_a \\ h_{41}x_a + h_{42}y_a + h_{43}z_a + h_{44}w_a \end{bmatrix} + \mu \begin{bmatrix} h_{11}x_b + h_{12}y_b + h_{13}z_b + h_{14}w_b \\ h_{21}x_b + h_{22}y_b + h_{23}z_b + h_{24}w_b \\ h_{31}x_b + h_{32}y_b + h_{33}z_b + h_{34}w_b \\ h_{41}x_b + h_{42}y_b + h_{43}z_b + h_{44}w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda \mathrm{HA} + \mu \mathrm{HB}$$

Matrix product

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\begin{bmatrix} h_{11}x_a + h_{12}y_a + h_{13}z_a + h_{14}w_a \\ h_{21}x_a + h_{22}y_a + h_{23}z_a + h_{24}w_a \\ h_{31}x_a + h_{32}y_a + h_{33}z_a + h_{34}w_a \\ h_{41}x_a + h_{42}y_a + h_{43}z_a + h_{44}w_a \end{bmatrix} + \mu\begin{bmatrix} h_{11}x_b + h_{12}y_b + h_{13}z_b + h_{14}w_b \\ h_{21}x_b + h_{22}y_b + h_{23}z_b + h_{24}w_b \\ h_{31}x_b + h_{32}y_b + h_{33}z_b + h_{34}w_b \\ h_{41}x_b + h_{42}y_b + h_{43}z_b + h_{44}w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\mathrm{HA} + \mu\mathrm{HB}$$

Definition

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\mathcal{H}(\mathcal{A}) + \mu\mathcal{H}(\mathcal{B})$$

# Homogeneous transformation

- Homogeneous transformation linearity

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\begin{bmatrix} h_{11}x_a + h_{12}y_a + h_{13}z_a + h_{14}w_a \\ h_{21}x_a + h_{22}y_a + h_{23}z_a + h_{24}w_a \\ h_{31}x_a + h_{32}y_a + h_{33}z_a + h_{34}w_a \\ h_{41}x_a + h_{42}y_a + h_{43}z_a + h_{44}w_a \end{bmatrix} + \mu\begin{bmatrix} h_{11}x_b + h_{12}y_b + h_{13}z_b + h_{14}w_b \\ h_{21}x_b + h_{22}y_b + h_{23}z_b + h_{24}w_b \\ h_{31}x_b + h_{32}y_b + h_{33}z_b + h_{34}w_b \\ h_{41}x_b + h_{42}y_b + h_{43}z_b + h_{44}w_b \end{bmatrix}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\mathrm{H}\mathcal{A} + \mu\mathrm{H}\mathcal{B}$$

$$\mathcal{H}(\lambda\mathcal{A}+\mu\mathcal{B}) = \lambda\mathcal{H}(\mathcal{A}) + \mu\mathcal{H}(\mathcal{B})$$

Any homogeneous transformation is a **linear application**

## Homogeneous Transformation

- Which kind of transformation can be represented ?
- Is it possible to use Homogeneous transformations to represents Euclidean transformations ?
- Is there transformation that can only be computed using Homogeneous coordinates ?

## Homogeneous Translation

- Let $\mathcal{X} = [x, y, z, w]^{\mathrm{T}}$ be a homogeneous vector. The translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$, denoted $\mathcal{T}(\alpha, \beta, \gamma)$, is such as:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X}$$

## Homogeneous Translation

- Let $\mathcal{X} = [x, y, z, w]^{\mathrm{T}}$ be a homogeneous vector. The translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$, denoted $\mathcal{T}(\alpha, \beta, \gamma)$, is such as:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X}$$

Where:

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Homogeneous Translation

- Let $\mathcal{X} = [x, y, z, w]^{\mathrm{T}}$ be a homogeneous vector. The translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$, denoted $\mathcal{T}(\alpha, \beta, \gamma)$, is such as:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

## Computing Euclidean translation

- Let $X = (x, y, z)$ be a vector within 3D Euclidean space.

## Computing Euclidean translation

- Let $X = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $X$ is: $\mathcal{X} = [x, y, z, 1]^{\mathsf{T}}$

## Computing Euclidean translation

- Let $X = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $X$ is: $\mathcal{X} = [x, y, z, 1]^{\mathsf{T}}$
- The homogeneous translation of $\mathcal{X}$ along $(\alpha, \beta, \gamma)$:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ 1 \end{bmatrix}$$

## Computing Euclidean translation

- Let $X = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $X$ is: $\mathcal{X} = [x, y, z, 1]^{\mathsf{T}}$
- The homogeneous translation of $\mathcal{X}$ along $(\alpha, \beta, \gamma)$:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ 1 \end{bmatrix}$$

- Euclidean representation: $[x + \alpha, y + \beta, z + \gamma, 1]^T \rightarrow \left( \frac{x + \alpha}{1}, \frac{y + \beta}{1}, \frac{z + \gamma}{1} \right) = (x + \alpha, y + \beta, z + \gamma)$

# Computing Euclidean translation

- Let $X = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $X$ is: $\mathcal{X} = [x, y, z, 1]^{\mathsf{T}}$
- The homogeneous translation of $\mathcal{X}$ along $(\alpha, \beta, \gamma)$:

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ 1 \end{bmatrix}$$

- Euclidean representation: $[x + \alpha, y + \beta, z + \gamma, 1]^T \rightarrow \left( \frac{x + \alpha}{1}, \frac{y + \beta}{1}, \frac{z + \gamma}{1} \right) = (x + \alpha, y + \beta, z + \gamma)$

$$\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) \rightarrow T(\alpha, \beta, \gamma)(X)$$

## Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

## Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ w \end{bmatrix}$$

# Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ w \end{bmatrix} = \begin{bmatrix} x + \alpha w - \alpha w \\ y + \beta w - \beta w \\ z + \gamma w - \gamma w \\ w \end{bmatrix}$$

## Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^\top$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ w \end{bmatrix} = \begin{bmatrix} x + \cancel{\alpha w} - \cancel{\alpha w} \\ y + \cancel{\beta w} - \cancel{\beta w} \\ z + \cancel{\gamma w} - \cancel{\gamma w} \\ w \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

# Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^\top$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x + \alpha \\ y + \beta \\ z + \gamma \\ w \end{bmatrix} = \begin{bmatrix} x + \cancel{\alpha w} - \cancel{\alpha w} \\ y + \cancel{\beta w} - \cancel{\beta w} \\ z + \cancel{\gamma w} - \cancel{\gamma w} \\ w \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \mathcal{X}$$

## Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \mathcal{X}$$

## Homogeneous translation inverse

- Let $\mathcal{X}$ and $\mathcal{Y}$ be a homogeneous vectors where $\mathcal{Y}$ is the result of the homogeneous translation of $\mathcal{X}$ along a vector $[\alpha, \beta, \gamma, 1]^{\mathrm{T}}$

$$\mathcal{X} = \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \qquad \mathcal{Y} = \mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X}) = \mathrm{T}\mathcal{X} = \begin{bmatrix} x + \alpha w \\ y + \beta w \\ z + \gamma w \\ w \end{bmatrix}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)(\mathcal{Y}) = T'\mathcal{Y} = \mathcal{X}$$

$$\mathcal{T}(-\alpha, -\beta, -\gamma)\big(\mathcal{T}(\alpha, \beta, \gamma)(\mathcal{X})\big) = T'T\mathcal{X} = \mathcal{X}$$

A Homogeneous translation $\mathcal{T}(\alpha, \beta, \gamma)$ is invertible and $\mathcal{T}(-\alpha, -\beta, -\gamma)$ is its inverse

# Homogeneous translation inverse

- Matrix inverse method

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{T}' = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous translation inverse

- Matrix inverse method

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{T}' = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{TT}' = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous translation inverse

- Matrix inverse method

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{T}' = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{TT}' = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -\alpha+\alpha \\ 0 & 1 & 0 & -\beta+\beta \\ 0 & 0 & 1 & -\gamma+\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous translation inverse

- Matrix inverse method

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{T}' = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{TT}' = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -\alpha+\alpha \\ 0 & 1 & 0 & -\beta+\beta \\ 0 & 0 & 1 & -\gamma+\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

# Homogeneous translation inverse

- Matrix inverse method

$$\mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{T}' = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{TT}' = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -\alpha+\alpha \\ 0 & 1 & 0 & -\beta+\beta \\ 0 & 0 & 1 & -\gamma+\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

$$\mathrm{TT}' = ID$$

T′ is the inverse matrix of T

# Homogeneous translation matrix structure

$$\mathrm{H} = \begin{bmatrix} 1 & 0 & 0 & \\ 0 & 1 & 0 & T \\ 0 & 0 & 1 & \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad T = \begin{bmatrix} t_1 \\ t_2 \\ t_3 \end{bmatrix}$$

# Homogeneous translation

- Definition

$$\mathcal{T}(\alpha,\beta,\gamma) = \mathrm{T} = \begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Properties
  - Linear application
  - Invertible: $\mathcal{T}^{-1}(\alpha,\beta,\gamma) = T^{-1} = \begin{bmatrix} 1 & 0 & 0 & -\alpha \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & -\gamma \\ 0 & 0 & 0 & 1 \end{bmatrix}$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^T$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\omega$ around $X$ axis, denoted $\mathcal{R}_x(\omega)$, is such as:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \mathrm{R}_x\mathcal{V}$$

Where:

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^T$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\omega$ around $X$ axis, denoted $\mathcal{R}_x(\omega)$, is such as:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \mathrm{R}_x\mathcal{V} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^T$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\omega$ around $X$ axis, denoted $\mathcal{R}_x(\omega)$, is such as:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \mathrm{R}_x\mathcal{V} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \begin{bmatrix} x \\ y\cos\omega - z\sin\omega \\ y\sin\omega + z\cos\omega \\ w \end{bmatrix}$$

## Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.

## Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^{\mathsf{T}}$

## Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \begin{bmatrix} x \\ y\cos\omega - z\sin\omega \\ y\sin\omega + z\cos\omega \\ 1 \end{bmatrix}$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \begin{bmatrix} x \\ y\cos\omega - z\sin\omega \\ y\sin\omega + z\cos\omega \\ 1 \end{bmatrix} \rightarrow \left(\frac{x}{1}, \frac{y\cos\omega - z\sin\omega}{1}, \frac{y\sin\omega + z\cos\omega}{1}\right)$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \begin{bmatrix} x \\ y\cos\omega - z\sin\omega \\ y\sin\omega + z\cos\omega \\ 1 \end{bmatrix} \rightarrow \left(\frac{x}{1}, \frac{y\cos\omega - z\sin\omega}{1}, \frac{y\sin\omega + z\cos\omega}{1}\right)$$

$$\rightarrow (x, y\cos\omega - z\sin\omega\,, y\sin\omega + z\cos\omega)$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_x(\omega)(\mathcal{V}) = \begin{bmatrix} x \\ y\cos\omega - z\sin\omega \\ y\sin\omega + z\cos\omega \\ 1 \end{bmatrix} \rightarrow \left(\frac{x}{1}, \frac{y\cos\omega - z\sin\omega}{1}, \frac{y\sin\omega + z\cos\omega}{1}\right)$$

$$\rightarrow (x, y\cos\omega - z\sin\omega, y\sin\omega + z\cos\omega)$$

$$\mathcal{R}_x(\omega)(\mathcal{V}) \rightarrow R_x(\omega)(V)$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}'_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(-\omega) & -\sin(-\omega) & 0 \\ 0 & \sin(-\omega) & \cos(-\omega) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}'_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(-\omega) & -\sin(-\omega) & 0 \\ 0 & \sin(-\omega) & \cos(-\omega) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\cos(-\omega) = \cos\omega$$

$$\mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin(-\omega) & 0 \\ 0 & \sin(-\omega) & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}'_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin(-\omega) & 0 \\ 0 & \sin(-\omega) & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\sin(-\omega) = -\sin\omega$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}'_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}'_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \mathbf{R}_x^{\top}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos^2\omega + \sin^2\omega & \cos\omega\sin\omega - \cos\omega\sin\omega & 0 \\ 0 & \sin\omega\cos\omega - \cos\omega\sin\omega & \sin^2\omega + \cos^2\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos^2\omega + \sin^2\omega & \cos\omega\sin\omega - \cos\omega\sin\omega & 0 \\ 0 & \sin\omega\cos\omega - \cos\omega\sin\omega & \sin^2\omega + \cos^2\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \cos\omega\sin\omega - \cos\omega\sin\omega & 0 \\ 0 & \sin\omega\cos\omega - \cos\omega\sin\omega & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\sin^2\omega + \cos^2\omega = 1$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \cos\omega\sin\omega - \cos\omega\sin\omega & 0 \\ 0 & \sin\omega\cos\omega - \cos\omega\sin\omega & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_x$ be the matrix of the homogeneous rotation $\mathcal{R}_x(\omega)$ and let $\mathrm{R}_x^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_x(-\omega)$

$$\mathrm{R}_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

A Homogeneous $X$ axis rotation $\mathcal{R}_x(\omega)$ is invertible and $\mathcal{R}_x(-\omega)$ is its inverse. Moreover, $\mathrm{R}_x^{-1} = \mathrm{R}_x^T$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^{\mathsf{T}}$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\varphi$ around $Y$ axis, denoted $\mathcal{R}_y(\varphi)$, is such as:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \mathrm{R}_y\mathcal{V}$$

Where:

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^{\mathsf{T}}$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\varphi$ around $Y$ axis, denoted $\mathcal{R}_y(\varphi)$, is such as:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \mathrm{R}_y\mathcal{V} = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^{\mathsf{T}}$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\varphi$ around $Y$ axis, denoted $\mathcal{R}_y(\varphi)$, is such as:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \mathrm{R}_y\mathcal{V} = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}$$

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \begin{bmatrix} x\cos\varphi + z\sin\varphi \\ y \\ z\cos\varphi - x\sin\varphi \\ w \end{bmatrix}$$

## Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.

## Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^{\mathsf{T}}$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $Y$ axis by an angle $\varphi$:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \begin{bmatrix} x\cos\varphi + z\sin\varphi \\ y \\ z\cos\varphi - x\sin\varphi \\ 1 \end{bmatrix}$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \begin{bmatrix} x\cos\varphi + z\sin\varphi \\ y \\ z\cos\varphi - x\sin\varphi \\ 1 \end{bmatrix} \rightarrow \left( \frac{x\cos\varphi + z\sin\varphi}{1}, \frac{y}{1}, \frac{z\cos\varphi - x\sin\varphi}{1} \right)$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \begin{bmatrix} x\cos\varphi + z\sin\varphi \\ y \\ z\cos\varphi - x\sin\varphi \\ 1 \end{bmatrix} \rightarrow \left( \frac{x\cos\varphi + z\sin\varphi}{1}, \frac{y}{1}, \frac{z\cos\varphi - x\sin\varphi}{1} \right)$$

$$\rightarrow (x\cos\varphi + z\sin\varphi\,, y, z\cos\varphi - x\sin\varphi)$$

# Computing Euclidean axis rotation

- Let $V = (x, y, z)$ be a vector within 3D Euclidean space.
- Homogeneous representation of $V$ is: $\mathcal{V} = [x, y, z, 1]^\top$
- Homogeneous rotation of $\mathcal{V}$ around $X$ axis by an angle $\omega$:

$$\mathcal{R}_y(\varphi)(\mathcal{V}) = \begin{bmatrix} x\cos\varphi + z\sin\varphi \\ y \\ z\cos\varphi - x\sin\varphi \\ 1 \end{bmatrix} \rightarrow \left( \frac{x\cos\varphi + z\sin\varphi}{1}, \frac{y}{1}, \frac{z\cos\varphi - x\sin\varphi}{1} \right)$$

$$\rightarrow (x\cos\varphi + z\sin\varphi\,, y, z\cos\varphi - x\sin\varphi)$$

$$\mathcal{R}_y(\varphi)(\mathcal{V}) \rightarrow R_y(\varphi)(V)$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}'_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_y = \begin{bmatrix} \cos(-\varphi) & 0 & \sin(-\varphi) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(-\varphi) & 0 & \cos(-\varphi) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}'_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_y = \begin{bmatrix} \cos(-\varphi) & 0 & \sin(-\varphi) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(-\varphi) & 0 & \cos(-\varphi) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_y = \begin{bmatrix} \cos\varphi & 0 & \sin(-\varphi) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(-\varphi) & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\cos(-\omega) = \cos\omega$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}'_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_y = \begin{bmatrix} \cos\varphi & 0 & \sin(-\varphi) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(-\varphi) & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}'_y = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\sin(-\omega) = -\sin\omega$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}'_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_y = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}'_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}'_y = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \mathbf{R}_y^\top$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} \cos^2\varphi + \sin^2\varphi & 0 & -\cos\varphi\sin\varphi + \sin\varphi\cos\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi\cos\varphi + \cos\varphi\sin\varphi & 0 & \cos^2\varphi + \sin^2\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} \cos^2\varphi + \sin^2\varphi & 0 & -\cos\varphi\sin\varphi + \sin\varphi\cos\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi\cos\varphi + \cos\varphi\sin\varphi & 0 & \cos^2\varphi + \sin^2\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} 1 & 0 & -\cos\varphi\sin\varphi + \sin\varphi\cos\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi\cos\varphi + \cos\varphi\sin\varphi & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\sin^2\omega + \cos^2\omega = 1$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} 1 & 0 & -\cos\varphi\sin\varphi + \sin\varphi\cos\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi\cos\varphi + \cos\varphi\sin\varphi & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

# Homogeneous rotation inverse

- Let $\mathrm{R}_y$ be the matrix of the homogeneous rotation $\mathcal{R}_y(\varphi)$ and let $\mathrm{R}_y^\top$ be the matrix of the homogeneous rotation $\mathcal{R}_y(-\varphi)$

$$\mathrm{R}_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R}_y^\top = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_y\mathrm{R}_y^\top = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = ID$$

A Homogeneous $Y$ axis rotation $\mathcal{R}_y(\varphi)$ is invertible and $\mathcal{R}_y(-\varphi)$ is its inverse. Moreover, $\mathrm{R}_y^{-1} = \mathrm{R}_y^T$

## Homogeneous axis rotation

- Let $\mathcal{V} = [x, y, z, w]^T$ be a homogeneous vector. The rotation of $\mathcal{V}$ by an angle $\kappa$ around $Z$ axis, denoted $\mathcal{R}_z(\kappa)$, is such as:

$$\mathcal{R}_z(\kappa)(\mathcal{V}) = \mathrm{R}_z\mathcal{V}$$

Where:

$$\mathrm{R}_z = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous axis rotation

- Transform Matrices

$$R_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad R_y = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad R_z = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Inverse transform matrices

$$R_x^{-1} = R_x^{\top} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad R_y^{-1} = R_y^{\top} = \begin{bmatrix} \cos\varphi & 0 & -\sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ \sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad R_z^{-1} = R_z^{\top} = \begin{bmatrix} \cos\varphi & \sin\kappa & 0 & 0 \\ -\sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Homogeneous rotation

- The rotation of angles $\omega, \varphi, \kappa$ respectively around $X, Y$ and $Z$ axis, denoted $\mathcal{R}_{xyz}(\omega, \varphi, \kappa)$, is such as:

$$\mathcal{R}_{xyz}(\omega, \varphi, \kappa) \equiv \mathrm{R}_z \mathrm{R}_y \mathrm{R}_x$$

# Homogeneous rotation

- The rotation of angles $\omega, \varphi, \kappa$ respectively around $X, Y$ and $Z$ axis, denoted $\mathcal{R}_{xyz}(\omega, \varphi, \kappa)$, is such as:

$$\mathcal{R}_{xyz}(\omega, \varphi, \kappa) \equiv \mathrm{R}_z \mathrm{R}_y \mathrm{R}_x$$

$$\mathrm{R}_z \mathrm{R}_y \mathrm{R}_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation

- The rotation of angles $\omega, \varphi, \kappa$ respectively around $X, Y$ and $Z$ axis, denoted $\mathcal{R}_{xyz}(\omega, \varphi, \kappa)$, is such as:

$$\mathcal{R}_{xyz}(\omega, \varphi, \kappa) \equiv \mathrm{R}_z \mathrm{R}_y \mathrm{R}_x$$

$$\mathrm{R}_z \mathrm{R}_y \mathrm{R}_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_z \mathrm{R}_y \mathrm{R}_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & \sin\varphi\sin\omega & \sin\varphi\cos\omega & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Homogeneous rotation

- The rotation of angles $\omega, \varphi, \kappa$ respectively around $X, Y$ and $Z$ axis, denoted $\mathcal{R}_{xyz}(\omega, \varphi, \kappa)$, is such as:

$$\mathcal{R}_{xyz}(\omega, \varphi, \kappa) \equiv \mathrm{R}_z \mathrm{R}_y \mathrm{R}_x$$

$$\mathrm{R}_z \mathrm{R}_y \mathrm{R}_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & \sin\varphi\sin\omega & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_z \mathrm{R}_y \mathrm{R}_x = \begin{bmatrix} \cos\kappa\cos\varphi & \cos\kappa\sin\varphi\sin\omega - \sin\kappa\cos\omega & \cos\kappa\sin\varphi\cos\omega + \sin\kappa\sin\omega & 0 \\ \sin\kappa\cos\varphi & \sin\kappa\sin\varphi\sin\omega + \cos\kappa\cos\omega & \sin\kappa\sin\varphi\cos\omega - \cos\kappa\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Homogeneous transform combination

- Homogeneous transforms combination is obtained by multiplying corresponding matrices.

- **Definition:** Let $\mathcal{F}_1, \dots, \mathcal{F}_n$ be transforms respectively represented by homogeneous $4 \times 4$ matrices $\mathrm{F}_1, \dots, \mathrm{F}_n$. The **composed transform** resulting from the composition of $\mathcal{F}_i$ transforms, defined by $\mathcal{G} = \mathcal{F}_1 \circ \cdots \circ \mathcal{F}_n$, is represented by the G matrix such as:

$$\mathrm{G} = \prod_{i=n}^{1} \mathrm{F}_i = \mathrm{F}_n \mathrm{F}_{n-1} \dots \mathrm{F}_1$$

# Rigid-body transformation

- **Definition:** A rigid body transformation $\mathcal{B}$ that combine translations and rotations is represented by a matrix B such as:

$$\mathrm{B} = \begin{bmatrix} & \mathrm{R} & & T \\ & & & \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

with $\mathrm{R} = \begin{bmatrix} r_{11} & r_{12} & r_{13} \\ r_{21} & r_{22} & r_{23} \\ r_{31} & r_{31} & r_{33} \end{bmatrix}$ and $\mathrm{T} = \begin{bmatrix} t_1 \\ t_2 \\ t_3 \end{bmatrix}$

## Rigid-body inverse transformation

- **Definition:** The inverse of a rigid body transformation $\mathcal{B}$ is a rigid body transformation $\mathcal{B}^{-1}$ that is represented by a matrix $\mathrm{B}^{-1}$ such as:

$$\mathrm{B}^{-1} = \begin{bmatrix} & \mathrm{R}^{-1} & & -\mathrm{R}^{-1}T \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} & \mathrm{R}^{\mathrm{T}} & & -\mathrm{R}^{\mathrm{T}}\mathrm{T} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\text{with } \mathrm{B} = \begin{bmatrix} & \mathrm{R} & & T \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Representing Transforms

| Transform | Computation | Linearity | Distances | Angles |
|---|---|---|---|---|
| Translation | $\begin{bmatrix} s_x r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & s_y r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & s_z r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ | Linear | ✓ | ✓ |
| Rotation | | Linear | ✓ | ✓ |
| Scale (uniform) | | Linear | ✗ | ✓ |
| Scale | | Linear | ✗ | ✗ |

$$\mathrm{B} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad \mathrm{R} = \begin{bmatrix} r_{11} & r_{12} & r_{13} \\ r_{21} & r_{22} & r_{23} \\ r_{31} & r_{32} & r_{33} \end{bmatrix} \qquad \mathrm{T} = \begin{bmatrix} t_1 \\ t_2 \\ t_3 \end{bmatrix} \qquad \mathrm{R}^\top = \begin{bmatrix} r_{11} & r_{21} & r_{31} \\ r_{12} & r_{22} & r_{32} \\ r_{13} & r_{23} & r_{33} \end{bmatrix}$$

$$-\mathrm{R}^\top\mathrm{T} = \begin{bmatrix} r_{11} & r_{21} & r_{31} \\ r_{12} & r_{22} & r_{32} \\ r_{13} & r_{23} & r_{33} \end{bmatrix} \begin{bmatrix} -r_{11}t_1 - r_{21}t_2 - r_{31}t_3 \\ -r_{12}t_1 - r_{22}t_2 - r_{32}t_3 \\ -r_{13}t_1 - r_{23}t_2 - r_{33}t_3 \end{bmatrix}$$

$$\mathrm{B}^{-1} = \begin{bmatrix} r_{11} & r_{21} & r_{31} & -r_{11}t_1 - r_{21}t_2 - r_{31}t_3 \\ r_{12} & r_{22} & r_{32} & -r_{12}t_1 - r_{22}t_2 - r_{32}t_3 \\ r_{13} & r_{23} & r_{33} & -r_{13}t_1 - r_{23}t_2 - r_{33}t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$V' = \begin{bmatrix} r_{11}v_1 + r_{12}v_2 + r_{13}v_3 + t_1 \\ r_{21}v_1 + r_{22}v_2 + r_{33}v_3 + t_2 \\ r_{31}v_1 + r_{32}v_2 + r_{33}v_3 + t_3 \\ 1 \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} r_{11}(r_{11}v_1 + r_{12}v_2 + r_{13}v_3 + t_1) + r_{12}(r_{21}v_1 + r_{22}v_2 + r_{33}v_3 + t_2) + r_{13}(r_{31}v_1 + r_{32}v_2 + r_{33}v_3 + t_1) + t_1 \\ \\ 0 \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} r_{11}r_{11}v_1 + r_{11}r_{12}v_2 + r_{11}r_{13}v_3 + r_{11}t_1 + r_{12}r_{21}v_1 + r_{12}r_{22}v_2 + r_{12}r_{33}v_3 + r_{12}t_2 + r_{13}r_{31}v_1 + r_{13}r_{32}v_2 + r_{13}r_{33}v_3 + r_{13}t_3 + t_1 \\ \\ 0 \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} r_{11}r_{11}v_1 + r_{12}r_{21}v_1 + r_{13}r_{31}v_1 + r_{11}r_{12}v_2 + r_{12}r_{22}v_2 + r_{13}r_{32}v_2 + r_{11}r_{13}v_3 + r_{13}r_{33}v_3 + r_{12}r_{33}v_3 + t_1r_{11} + t_1 + r_{12}t_2 + r_{13}t_3 \\ \\ 0 \end{bmatrix}$$

$$\mathrm{V} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ 1 \end{bmatrix} \quad \mathrm{BV} = \begin{bmatrix} r_{11}v_1 + r_{12}v_2 + r_{13}v_3 + t_1 \\ r_{21}v_1 + r_{22}v_2 + r_{33}v_3 + t_2 \\ r_{31}v_1 + r_{32}v_2 + r_{33}v_3 + t_1 \\ 1 \end{bmatrix} = \mathrm{V}'$$

$$\mathrm{B}^{-1} = \begin{bmatrix} r_{11} & r_{21} & r_{31} & -r_{11}t_1 - r_{21}t_2 - r_{31}t_3 \\ r_{12} & r_{22} & r_{32} & -r_{12}t_1 - r_{22}t_2 - r_{32}t_3 \\ r_{13} & r_{23} & r_{33} & -r_{13}t_1 - r_{23}t_2 - r_{33}t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{B}^{-1} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} r_{11}^2 + r_{12}^2 + r_{13}^2 & & & \\ & & & \\ & & & \\ 0 & & 0 \quad 0 & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} kx_a + jx_b \\ ky_a + jy_b \\ kz_a + jz_b \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} kx_a + jx_b + \alpha(k+j) \\ ky_a + jy_b + \beta(k+j) \\ kz_a + jz_b + \gamma(k+j) \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} kx_a + jx_b \\ ky_a + jy_b \\ kz_a + jz_b \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} kx_a + jx_b + \alpha k + \alpha j \\ ky_a + jy_b + \beta k + \beta j \\ kz_a + jz_b + \gamma k + \gamma j \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} kx_a + jx_b \\ ky_a + jy_b \\ kz_a + jz_b \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} kx_a + \alpha k + jx_b + \alpha j \\ ky_a + \beta k + jy_b + \beta j \\ kz_a + \gamma k + jz_b + \gamma j \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} kx_a + jx_b \\ ky_a + jy_b \\ kz_a + jz_b \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} kx_a + \alpha k \\ ky_a + \beta k \\ kz_a + \gamma k \\ kw_a \end{bmatrix} + \begin{bmatrix} jx_b + \alpha j \\ jy_b + \beta j \\ jz_b + \gamma j \\ jw_b \end{bmatrix}$$

$$\begin{bmatrix} kx_a + jx_b \\ ky_a + jy_b \\ kz_a + jz_b \\ kw_a + jw_b \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & \alpha \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & \gamma \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad k \begin{bmatrix} x_a + \alpha \\ y_a + \beta \\ z_a + \gamma \\ w_a \end{bmatrix} + j \begin{bmatrix} jx_b + \alpha \\ jy_b + \beta \\ jz_b + \gamma \\ w_b \end{bmatrix}$$

$$\mathrm{R}_x\mathrm{R}_x^T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & \sin\omega & 0 \\ 0 & -\sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos^2\omega + \sin^2\omega & \cos\omega\sin\omega - \cos\omega\sin\omega & 0 \\ 0 & \sin\omega\cos\omega - \cos\omega\sin\omega & \sin^2\omega + \cos^2\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{31} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} r_{11}x + r_{12}y + r_{13}z + t_1w \\ r_{21}x + r_{22}y + r_{23}z + t_2w \\ r_{31}x + r_{32}y + r_{33}z + t_3w \\ w \end{bmatrix}$$

$$\begin{bmatrix} r_{11} & r_{12} & r_{13} & 0 \\ r_{21} & r_{22} & r_{23} & 0 \\ r_{31} & r_{31} & r_{33} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & t_1 \\ 0 & 1 & 0 & t_2 \\ 0 & 0 & 1 & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & r_{11}t_1 + r_{12}t_2 + r_{13}t_3 \\ r_{21} & r_{22} & r_{23} & r_{21}t_1 + r_{22}t_2 + r_{23}t_3 \\ r_{31} & r_{32} & r_{33} & r_{31}t_1 + r_{32}t_2 + r_{33}t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & t_1 \\ 0 & 1 & 0 & t_2 \\ 0 & 0 & 1 & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} & r_{13} & 0 \\ r_{21} & r_{22} & r_{23} & 0 \\ r_{31} & r_{31} & r_{33} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_1 \\ r_{21} & r_{22} & r_{23} & t_2 \\ r_{31} & r_{32} & r_{33} & t_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_z R_y R_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & \sin\varphi\sin\omega & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_z R_y R_x = \begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\kappa\cos\varphi & \cos\kappa\sin\varphi\sin\omega - \sin\kappa\cos\omega & \sin\kappa\sin\omega & 0 \\ \sin\kappa\cos\varphi & \sin\kappa\sin\varphi\sin\omega + \cos\kappa\cos\omega & -\sin\kappa\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\kappa\cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_z R_y R_x = \begin{bmatrix} \cos\kappa\cos\varphi & \sin\varphi\sin\omega & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ 0 & \sin\omega & \cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathrm{R}_z\mathrm{R}_y\mathrm{R}_x = \begin{bmatrix} \cos\varphi & 0 & \sin\varphi & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\varphi & 0 & \cos\varphi & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\varphi & \sin\varphi\sin\omega & \sin\varphi\cos\omega & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$(x, y, z, w) = (2x, 2y, 2z, 2w)$

$(x, y, z, w) = (x/w, y/w, z/w, 1)$

$(x, y, z, w) = (1/x, 1/y, 1/z, 1/w)$

$(x, y, z, w) = (\alpha x, \alpha y, \alpha z, \alpha w), \alpha \in \mathbb{R}$

$$R_z R_y R_x = \begin{bmatrix} \cos\kappa\cos\varphi & \cos\kappa\sin\varphi\sin\omega - \sin\kappa\cos\omega & \cos\kappa\sin\varphi\cos\omega + \sin\kappa\sin\omega & 0 \\ \sin\kappa\cos\varphi & \sin\kappa\sin\varphi\sin\omega + \cos\kappa\cos\omega & \sin\kappa\sin\varphi\cos\omega - \cos\kappa\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} \cos\varphi & \sin\varphi\sin\omega & \sin\varphi\cos\omega & 0 \\ 0 & \cos\omega & -\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} \cos\kappa & -\sin\kappa & 0 & 0 \\ \sin\kappa & \cos\kappa & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos\kappa\cos\varphi & \cos\kappa\sin\varphi\sin\omega - \sin\kappa\cos\omega & \cos\kappa\sin\varphi\cos\omega + \sin\kappa\sin\omega & 0 \\ \sin\kappa\cos\varphi & \sin\kappa\sin\varphi\sin\omega + \cos\kappa\cos\omega & \sin\kappa\sin\varphi\cos\omega - \cos\kappa\sin\omega & 0 \\ -\sin\varphi & \cos\varphi\sin\omega & \cos\varphi\cos\omega & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$H^{-1} = \begin{bmatrix} -R & -T \\ 0\ 0\ 0 & 1 \end{bmatrix}$$

$$H^{-1} = \begin{bmatrix} R^{T} & -R^{T}T \\ 0\ 0\ 0 & 1 \end{bmatrix}$$

$$H^{-1} = \begin{bmatrix} R^{-1} & T^{-1} \\ 0\ 0\ 0 & 1 \end{bmatrix}$$

$$H^{-1} = \begin{bmatrix} -R & -RT \\ 0\ 0\ 0 & 1 \end{bmatrix}$$

$$H^{-1} = \begin{bmatrix} -R & -RT \\ 0\ 0\ 0 & 1 \end{bmatrix}$$